

7. Nickell, L. G. Embryo culture of weeping crabapple. Proc. Amer. Soc. Hort. Sci. 57: 401-405. 1951.
8. Nickell, L. G. Aseptic studies on metabolism of nitrogenous compounds in plants. Bull. Torrey Bot. Club 83: 421-427. 1956.
9. Nickell, L. G., and A. C. Finlay. Antibiotics and their effects on plant growth. Journ. Agr. and Food Chem. 2: 178-182. 1954.
10. Svenson, H. K. The new world species of *Azolla*. Amer. Fern Journ. 34: 69-84. 1944.
11. Wetmore, R. H., and G. Morel. Growth and development of *Adiantum pedatum* L. on nutrient agar. Amer. Journ. Bot. 36: 805-806. 1949 (Abstract).
12. Wetmore, R. H., and G. Morel. Polyphenol oxidase as a problem in organ culture and auxin diffusion studies of horsetails and ferns. Amer. Journ. Bot. 36: 830. 1949. (Abstract).
13. Wetmore, R. H., and C. Pratt. The growth and auxin relations of leaves of the maidenhair fern, *Adiantum pedatum* L. Amer. Journ. Bot. 36: 806. 1949. (Abstract).

A New Jamaican Species of *Ctenitis*

GEORGE R. PROCTOR

During the past several years the writer has explored some of the more remote corners of Jamaica for ferns, as a result of which a number of interesting new records have come to light. The type locality of the present new species is a precipitous mountain of white limestone called Dolphin Head, near the western end of the island in the Parish of Hanover. Here it grows among moist shaded boulders in association with other plants endemic to the area.

CTENITIS dolphinensis Proctor, *sp. nov.*

Ctenitis sect. *Hirtae* rhizomate crasso, suberecto, paleis aureo-brunneis, linearibus vel lineari-lanceolatis, attenuatis, integris dense vestito; stipitibus fasciculatis, brunneis, 30-35 cm. longis, ubique—versus basim densius—paleis bicoloribus filiformibus basi expansis vestitis;